

8296

L. C. BROWN

1916

21

Connecticut
FORMICOIDEA

By
WILLIAM MORTON WHEELER

Reprinted from Bulletin 22
of the
Connecticut Natural History and Geological Survey

1916, pp. 577-601.

FORMICOIDEA.

FORMICIDÆ.

By WILLIAM MORTON WHEELER.

The ants (family Formicidæ) are social Hymenopterous insects, and may be distinguished from the social bees and wasps by having workers, or neuters, as they are less appropriately called, without wings. They are, moreover, readily distinguished from these and all other Hymenopterous insects by the following characters:

1. The first antennal joint in the workers and females, and often also in the males, is greatly elongated and forms what is known as the scape. The remaining shorter joints, constituting the funiculus, or flagellum, are articulated at an angle with the scape and can be folded up against it.

2. One or two of the segments of the base of the abdomen are much reduced in size to form a pedicel, and these segments are either nodiform or bear an erect or inclined scale. When only one of these segments is present, it is known as the petiole; when two are present, the first is the petiole, the second the post-petiole. The swollen portion of the abdomen behind the pedicel is known as the gaster, and has one more visible segment in the male than in the female (queen).

3. The legs of ants are distinguished from those of many other Hymenoptera in having only one instead of two small joints (trochanters) between the hip (coxa) and femur.

4. The venation of the wings of male and female ants is much simplified and differs considerably from that of other Hymenoptera. The female, or queen ant, unlike the queens of the social bees and wasps, loses her wings after fertilization.

The colonies of all our northern ants nest either in the ground or in decaying wood. The nests, or formicaries, may be under stones or logs, and always consist of irregularly excavated, intercommunicating cavities, unlike the regular paper or waxen combs of other social Hymenoptera. Often the nests are surmounted

by earthen craters or dome-shaped mounds, or "hills." The latter are perforated with cavities which serve as incubators for the young, that is, for the minute eggs, the legless, grub-like larvæ, and the pupæ. The pupæ are either naked or enclosed in elliptical cocoons which are spun by the mature larvæ.

Many species of ants harbor in their nests messmates or parasites belonging to various groups of insects. Some of these so-called myrmecophiles are fed and cared for by the ants, others prey upon the ants or their brood. Certain species of ants may themselves become parasitic on other ants. A few of these parasitic species have lost their worker caste completely, and are, therefore, represented only by male and female individuals like the non-social Hymenoptera.

The food of ants consists primarily of other insects found dead or in a moribund or helpless condition on the ground or vegetation. Many species, however, feed on honey-dew, and either collect this sweet liquid directly from the plant-lice and scale insects of which it is the excrement, or lap it up from the surfaces of the leaves on which it has fallen. Ants are, on the whole, beneficial insects, since they consume enormous numbers of dead and decomposing organisms. Many of the less abundant species are neither beneficial nor noxious. A few, like the little red house-ant (*Monomorium pharaonis*) and the large black carpenter-ant (*Camponotus pennsylvanicus*), are sometimes a pest in houses. Both of these species are very fond of feeding on sweets in pantries, kitchens, etc., and the carpenter-ant also has the injurious habit of excavating its galleries in the beams and rafters of houses. A few species, like the garden ant (*Lasius americanus*) and the silky ant (*Formica subsericea*), disfigure lawns and garden beds with their burrows and craters.

The following list of ants occurring in Connecticut has been prepared at the suggestion of Dr. W. E. Britton from material collected by himself, Mr. H. L. Viereck, and others in various parts of the state, and from my own collections made during several summers in the vicinity of Colebrook, Winsted, and Norfolk in the Litchfield Hills. This list is probably very incomplete, as I have found several species in adjacent portions of New York (*e. g.*, near White Plains), not represented in the material from Connecticut. Previous authors have recorded from the

latter state several species which I have had to discard. Mayr ascribed to Connecticut *Pogonomyrmex subdentatus*, an ant known to occur only in the arid deserts of the Southwest; and Buckley described the following species from the same state: *Formica nortonii*, *F. americana*, *F. connecticutensis*, *F. gnava*, *F. occidentalis*, and *Myrmica (Diplorhoptrum) scabrata*. With the exception of *F. gnava*, none of these forms can be recognized from Buckley's abominable descriptions. Under *F. gnava* he evidently included several different ants. One of these, a form of *F. fusca* intermediate between the varieties *subsericea* Say and *neorufibarbis* Emery, I have been able to recognize in the Texan fauna, and I have therefore restricted Buckley's name to this particular variety. With this single exception, however, all of the above names of Buckley's Formicidæ may be consigned to oblivion.

As the worker caste is the best known and most commonly met with, it is the only one used for identification in the tables published in the following pages. These tables include the subfamilies, genera, and subgenera known to occur in North America north of Mexico. Of the five subfamilies only four are represented in the Northern States, the remaining one (Dorylinæ) being confined to tropical and subtropical regions.

Key to Subfamilies.

1. Cloacal orifice ventral, slit-shaped; sting well developed or vestigial; abdominal pedicel consisting of one or two segments 2
 Cloacal orifice terminal, circular, surrounded by a fringe of hairs; abdominal pedicel consisting of only a single segment; no constriction between the first and second gastric segments; pupæ usually enclosed in cocoon
CAMPONOTINÆ p. 590
2. Sting developed, sometimes very small but nevertheless exertile; abdominal pedicel consisting of one or two segments; when of only one, a distinct constriction between first and second gastric segments 3
 Sting vestigial; abdominal pedicel consisting of a single segment; no constriction between first and second gastric segments; anal glands which produce a secretion with a peculiar rancid-butter odor ("Tapinoma odor") often present; pupæ naked
DOLICHODERINÆ p. 589
3. Pupæ always enclosed in cocoons; abdominal pedicel consisting of a single segment; gaster with a distinct constriction between its first and second segments; frontal carinæ

- separated or close together; when close together, dilated to form oblique or horizontal laminae partly covering insertions of antennaePONERINÆ p. 580
- Pupæ naked; abdominal pedicel consisting of two segments 4
4. Frontal carinae very close together, almost vertical, not at all covering antennal insertions; eyes always very small or absent; tropical and subtropicalDORYLINÆ
- Frontal carinae of a different conformation and covering the antennal insertions; eyes rarely vestigial or absent; cosmopolitanMYRMICINÆ p. 581

PONERINÆ.

Key to Genera.

1. Frontal carinae closely approximated; antennae inserted very near oral margin; tip of gaster strongly deflected downward 2
- Frontal carinae of a different conformation; tip of gaster not deflected downward 3
2. Front of clypeus projecting in middle; petiole nodiform
Sysphincta
Clypeus not projecting in middle; petiole surmounted by a scale**Proceratium**
3. Mandibles linear, inserted close together at middle of oral border; petiole terminating in a point or spine above
Odontomachus
Mandibles inserted at corners of head; petiole rounded or flattened above 4
4. Antennae very thick and robust 5
- Antennae not greatly thickened 6
5. Pygidium with a row of prominent prickles on its lateral border; last antennal joint not greatly enlarged
Acanthostichus
Pygidium without prominent prickles on its lateral border; last antennal joint greatly enlarged**Cerapachys**
6. Mandibles long and slender, with coarse, bidenticulate teeth; clypeus with numerous teeth on its anterior border; petiole not constricted posteriorly**Stigmatomma** p. 581
- Of a different conformation 7
7. Claws pectinate 8
- Claws simple 9
8. Mandibles edentate, slender; without distinct apical border
Leptogenys (s. str.)
Mandibles broader, generally toothed; with distinct apical border**Leptogenys (Lobopelta)**
9. Median spur of mid and hind legs alone developed; lateral spurs lacking; small species with vestigial eyes..**Ponera** p. 581

- Both spurs of mid and hind legs well developed; medium
or large species, with larger eyes 10
10. Cheeks with a longitudinal carina **Neoponera**
Cheeks without a carina 11
11. Pronotum more or less marginate on sides; mid tibiæ not
abbreviated nor beset with prominent bristles
Pachycondyla (*s. str.*)
Pronotum not marginate on sides; mid tibiæ short, with
prominent bristles on their exterior surfaces
Euponera (**Pseudoponera**)

Stigmatomma Roger.

S. pallipes Haldeman, var **wheeleri** Santschi.

This singular and primitive ant is subterranean or hypogæic in its habits, and occurs only in rich, rather damp woods, under stones, leaf-mold, or more rarely under rotten logs. It is by no means common. The colonies are small, comprising in extreme cases from forty to sixty individuals. The males and winged females appear in August and early September.

Suffield (Geo. Dimmock); Colebrook (W. M. W.).

Ponera Latreille.

P. coarctata pennsylvanica (Buckley) Emery.

Like the preceding, this small, slender species lives in small colonies, but is much more abundant. It nests under stones and vegetable mold, in rotten logs, etc., in rather open woods, along hedges, etc. The males and winged females appear in late August and early September.

Colebrook (W. M. W.).

MYRMICINÆ.

Key to Genera.

1. Workers absent **Epæcus**; **Sympheidole**; **Epipheidole**
Workers present 2
2. Clypeus not extending back between frontal carinæ, which
are closely approximated; antennæ 12-jointed
Pseudomyrma
Clypeus almost always extending back between frontal carinæ,
which are more or less separated; in the opposite case
antennæ 11-jointed 3
3. Antennal fossæ prolonged as grooves for antennal scapes

along sides of head dorsal to eyes and covered by extended lateral margins of head; antennæ 11-jointed

Cryptocerus

- Antennal fossæ of a different conformation or antennæ of a different number of joints 4
4. Postpetiole articulated to dorsal surface of gaster, which is flattened dorsally, more convex ventrally, and pointed at tip **Crematogaster** p. 585
- Postpetiole inserted at anterior end of gaster, which is of the usual shape 5
5. Antennæ 6-jointed; head cordiform, antennal fossæ as long as scapes **Strumigenys**
- Antennæ with more than six joints 6
6. Antennæ 11-jointed; without a distinct club or with a club consisting of only a single joint 7
- Antennal club consisting of several joints, or antennæ not 11-jointed 10
7. Integument rough, bearing stiff or hooked hairs 8
- Integument smoother; hairs scale-like and appressed **Cyphomyrmex**
8. Large species; workers highly polymorphic; head with a pair of occipital spines only; thorax with three pairs of dorsal spines or tubercles **Atta** (*s. str.*)
- Small species; workers monomorphic or feebly polymorphic; thoracic dorsum with four pairs of spines or tubercles 9
9. Head broad with rounded occipital lobes, without supraocular spines or tubercles **Atta** (**Moellerius**)
- Head narrow, with angular occipital lobes; body rough, covered with small tubercles **Atta** (**Trachymyrmex**)
10. Antennæ with a 2-jointed club 11
- Antennal club, when developed, with more than two joints 12
11. Antennæ 10-jointed, epinotum unarmed..... **Solenopsis** p. 584
- Antennæ 11-jointed, epinotum dentate **Erebomyrma**
12. Posterior margin of clypeus elevated in the form of a welt or ridge bordering antennal fossa in front 13
- Posterior border of clypeus not thus elevated 15
13. Portion of clypeus in front of antennal insertion narrow but not reduced to a mere ridge; antennæ of male 10-jointed.. 14
- Portion of clypeus in front of antennal insertion reduced to a mere ridge; antennæ of male 13-jointed .. **Myrmecina** p. 584
14. Antennæ 12-jointed **Tetramorium** (*s. str.*)
- Antennæ 11-jointed **Tetramorium** (**Xiphomyrmex**)
15. Antennæ 11-jointed 16
- Antennæ 12-jointed 19

16. Thorax and petiole without any traces of teeth or spines; pronotum never angular 17
 Epinotum armed with spines or teeth 18
17. Petiole distinctly pedunculate **Monomorium** p. 584
 Petiole not pedunculate **Xenomymex**
18. Mesoëpinotal constriction distinct; males ergatomorphic
Symmyrmica
 Mesoëpinotal constriction faint or lacking; males not ergatomorphic **Leptothorax** p. 588
19. Workers strongly dimorphic, usually without intermediates connecting the extreme forms; antennal club 3-jointed, longer than remainder of funiculus **Pheidole** p. 584
 Workers monomorphic or polymorphic, *i. e.*, with mediæ intermediate between major and minor forms; antennal club indistinct or shorter than remainder of funiculus 20
20. Last three antennal joints much shorter than remainder of funiculus and not forming a distinct club 21
 Last three antennal joints forming a distinct club nearly as long as remainder of funiculus 26
21. Thoracic dorsum impressed at mesoëpinotal suture; promeso-notal suture usually distinct 22
 Thoracic dorsum without any traces of suture or impressions
Pogonomyrmex
22. Posterior tibial spurs pectinated **Myrmica** p. 587
 Posterior tibial spurs simple 23
23. Small hypogæic species, with vestigial eyes and two keels on clypeus **Stenammas** p. 585
 Medium-sized epigæic species with well-developed eyes and no keels on clypeus 24
24. Workers monomorphic 25
 Workers polymorphic **Novomessor**
25. Cosmopolitan species with moderately slender thorax and legs **Aphænogaster (s. str.)** p. 585
 Tropical and subtropical species with very slender thorax and legs **Aphænogaster (Deromyrma)**
26. Clypeus armed with a pair of ridges which project forward in the form of teeth, rarely without teeth, but then the epinotum quite unarmed; mesoëpinotal suture marked
Monomorium p. 584
 Clypeus of a different conformation, rarely 2-toothed, but then the mesoëpinotal suture indistinct 27
27. Postpetiole campanulate, not constricted behind, but applied with its whole posterior surface to first gastric segment
Macromischa
 Postpetiole constricted behind **Leptothorax** p. 588

Myrmecina Curtis.

M. graminicola americana var. **brevispinosa** Emery.

Rare; nesting in small colonies under stones in shady woods. Males and winged females appear during August. It is a timid species which "feigns death" when rudely handled.

Colebrook (W. M. W.).

Monomorium Mayr.

M. minimum (Buckley) Emery.

This very small jet-black ant nests in small crater nests in sandy or gravelly places. The workers move in files, visiting plants in search of honey-dew and the secretion of the extrafloral nectaries. The species seems to be absent from the hilly portions of the State.

New Haven, North Haven (H. L. V.).

°**M. pharaonsis** Linnæus.

This little "red" or "yellow house ant," though not recorded from Connecticut, can hardly be absent from the seaport towns, as it is common on ships and has been carried to all parts of the world from its original home in the warmer regions of the Old World.

Solenopsis Westwood.

S. molesta Say.

A species with minute yellow workers and much larger brown females and blackish males. It is common in open grassy places, where it may live either in independent formicaries under stones, or as a thief ant in the walls separating the galleres of the formicaries of larger ants belonging to the genera *Formica*, *Myrmica*, *Aphaenogaster*, etc. The males and winged females appear late in August.

New Haven (E. B. Whittlesey); North Haven (H. L. V.); Colebrook (W. M. W.).

Pheidole Westwood.

P. pilifera Roger.

This ant undoubtedly occurs in sandy regions in the southern portion of the State, as it is common on Long Island (Cold Spring Harbor) and has been found in Massachusetts. It is a true harvesting ant, storing the chambers of its nest with seeds of grass

and other plants. The huge-headed soldiers undoubtedly function as seed-crushers.

New Haven (W. E. B.).

Crematogaster Lund.

C. lineolata Say.

A very common species, nesting under stones in open places, under stumps, boards, the bark of old logs, etc. There is a vestigial tendency in this ant to construct carton partitions or cells in its nest or over aphids and coccids on plants. The workers, which have a disagreeable odor, move about in loose files and often carry the triangular gaster over the thorax with the tip turned forward. The males and winged females may be found in the nests from the latter part of July to September.

Connecticut (Mayr); Branford, West Haven (H. L. V.); New Haven, New Canaan (W. E. B.); Suffield (Geo. Dimmock); Colebrook (W. M. W.).

C. lineolata var. *cerasi* Fitch.

Differs from the preceding in its paler color.

Colebrook (W. M. W.).

Stenamamma Mayr.

S. brevicorne Mayr.

Rare; nesting in small colonies under stones or vegetable mold in rich woods.

Colebrook (W. M. W.).

Aphænogaster Mayr.

Key to Species.

1. Antennal scape with a long, flattened lobe at its base *treatæ*
Antennal scape without a lobe 2
2. Basal third of first gastric segment longitudinally striated
mariae
Basal third of first gastric segment smooth 3
3. Epinotal spines at least as long as base of epinotum; color
red *tennesseensis*
Epinotal spines shorter than base of epinotum; color red-
dish brown or black 4
4. Epinotal spines somewhat longer than half the base of epi-
notum; length 4.5-5 mm. (typical) *fulva*
Epinotal spines shorter; length 4-4.5 mm. 5

5. Color reddish brown *fulva* subspecies *aquia*
 Color pitchy black *fulva aquia* var. *picea*

A. tennesseensis Mayr.

This species differs from our other species of *Aphænogaster* in having very small and very smooth females with huge epinotal spines. These aberrant females probably establish their colonies in nests of *Aphænogaster fulva* or some one of its varieties, in the same way that *Formica difficilis* var. *consocians* establishes its colonies in nests of *F. schaufussi* var. *incerta* (*vide infra*). At least *tennesseensis* is known to occur only in regions where *fulva* is unusually abundant, and several mixed colonies of the two species, containing queens of *tennesseensis* only, have been recorded. When living in unmixed colonies it always nests in rotten wood.

Colebrook (W. M. W.).

A. treatæ Forel.

The female and worker are easily recognized by the remarkable lamella on the base of the antennal scape.

Poquonock (H. L. V.), almost the northernmost locality in which this species has been found.

A. mariæ Forel.

A single winged female that had just descended from her nuptial flight was taken 8 September, 1901.

Colebrook (W. M. W.).

A. fulva Roger.

Nesting in rotten wood in rather dense forests; rarer than the following subspecies and variety.

Connecticut (Mayr); Colebrook (W. M. W.).

A. fulva aquia (Buckley) Emery.

Under stones in shady woods, often in the same stations as the following variety.

Branford (H. L. V., H. W. W.); New Haven (H. L. V.); Colebrook (W. M. W.).

A. fulva aquia var. *picea* Emery.

Apparently common throughout the State. The males and winged females appear during July and August.

Connecticut (Emery); Colebrook (W. M. W.).

Myrmica Latreille.*Key to Species.*

1. First gastric segment with coarse, scattered punctures
punctiventris
- First gastric segment without such punctures 2
2. Antennal scape not dilated to form a tooth or lobe at base,
but merely curved **brevinodis** and varieties
- Antennal scape toothed or lobed at base
scabrinodis and varieties

M. punctiventris Roger.

A rare species nesting in small colonies under stones or moss in moist shady woods. It is easily recognized by the coarse punctures on the gaster of the worker and female. The winged phases appear during August and September.

Colebrook (W. M. W.).

M. brevinodis Emery, var. *canadensis* Wheeler.

In Connecticut this form is confined to the bogs and low-lying pastures among the Litchfield Hills where it nests in grassy hummocks or under stones. It is the host of a species of *Leptothorax*, *L. emersoni* (see p. 588). The males and winged females appear during August.

Colebrook (W. M. W.).

M. scabrinodis Nylander, var. *sabuleti* Meinert.

This variety of the palearctic *scabrinodis* is reddish in color and in the male phase has the antennal scape somewhat more than a third the length of the funiculus. It nests in sandy or gravelly, sunny places, such as open pastures, roadsides, etc. The males and winged females may be found in the nests in the latter part of August.

West Haven, Branford (H. L. V.); New Haven (W. E. B.);
Colebrook (W. M. W.).

M. scabrinodis var. *schencki* Emery.

This form sometimes passes in the literature as *lobicornis*. The male has short, thick antennal scapes, shorter than those of *sabuleti* and rarely longer than one-fourth of the funiculus.

Stafford (W. E. B.); Colebrook (W. M. W.).

M. scabrinodis var. *fracticornis* Emery.

A form which is occasionally found nesting in the grass of

L. curvispinosus Mayr, subspecies **ambiguus** Emery.

Very similar to the preceding but with shorter and straighter epinotal spines.

West Haven (H. L. V.); Stafford (W. E. B.); Colebrook (W. M. W.).

Tetramorium Mayr.°**T. cæspitum** Linnæus.

Though this form has not yet been recorded from Connecticut, there can be little doubt that it occurs within the state. I have found it at Mamaroneck and Cold Spring Harbor, N. Y., both localities very near the Connecticut boundary. It has been introduced into America from Europe.

DOLICHODERINÆ.*Key to Genera.*

1. Chitinous integument hard and brittle, often strongly sculptured; thorax and petiole often spinose or angular **Dolichoderus** p. 589
 Chitinous integument thin and flexible, smooth or very finely sculptured; thorax and petiole always unarmed 2
2. Scale of petiole very small, strongly inclined forward, or even altogether absent 3
 Scale of petiole more or less inclined, but well developed 4
3. Scale of petiole small but distinct; gizzard with a convex, 4-lobed calyx **Forelius**
 Scale vestigial or absent; gizzard with a depressed calyx, without lobes **Tapinoma** p. 590
4. Epinotum with a conical elevation **Dorymyrmex**
 Epinotum without a conical elevation 5
5. Body not conspicuously hairy or pubescent; gizzard very short with a large reflected calyx; ocelli absent .. **Iridomyrmex**
 Body densely pubescent; gizzard at least as long as broad; ocelli usually present in large workers **Liometopum**

Dolichoderus Lund.**D. mariæ** Forel.

Readily distinguished from our other species of *Dolichoderus* by the bright red head and thorax in the worker and female. It forms large colonies, nesting in sandy places about the roots of grasses and bushes. The workers ascend trees in files and attend aphids and coccids.

Connecticut (Emery).

D. plagiatus Mayr.

The head and thorax of the worker are coarsely punctate or foveolate and the gaster has large yellowish red spots. It nests in the ground in small colonies. In other respects its habits resemble those of the preceding species.

Rockville (H. L. V.); Colebrook (W. M. W.).

Tapinoma Foerster.**T. sessile** Say.

Evidently very common, especially in the southern portion of the state. It nests under stones, dead leaves, logs, bark, etc., usually in sunny places. The larvæ and pupæ are salmon-colored. The workers emit a peculiar rancid-butter odor, the characteristic "Tapinoma odor," which serves to distinguish them from all our other eastern ants.

Branford, New Haven, Stony Creek, Double Beach (H. L. V.); Orange (W. E. B.); Colebrook (W. M. W.).

CAMPONOTINÆ.

Key to Genera.

1. Antennæ 9-jointed **Brachymyrmex** p. 591
Antennæ with more than nine joints 2
2. Workers strongly polymorphic, *i. e.*, with large-headed workers (majores) and small-headed workers (minores) and intermediate forms (mediæ) **Camponotus** p. 600
Workers not polymorphic though often of variable size 3
3. Clypeal fossa distinctly separated from antennal fossa 4
Clypeal fossa confluent with antennal fossa 5
4. Antennal scapes and tibiæ with erect hairs; mesonotum constricted but not subcylindrical. **Prenolepis (Nylanderia)** p. 591
Antennal scapes and tibiæ without erect hairs; mesonotum strongly constricted and subcylindrical
Prenolepis (s. str.) p. 591
5. Second to fifth joints of funiculus shorter or not longer than succeeding joints; ocelli usually absent 6
Second to fifth joints of funiculus longer than succeeding joints; ocelli distinct 7
6. Maxillary palpi 6-jointed **Lasius (s. str.)** p. 591
Maxillary palpi 3-jointed **Lasius (Acanthomyops)** p. 594
7. Fourth joint of maxillary palpi nearly as long as fifth
Myrmecocystus
Fourth joint of maxillary palpi a little longer than fifth 8
8. Mandibles with broad dentate apical border **Formica** p. 594
Mandibles narrow, falcate and pointed **Polyergus** p. 599

Brachymyrmex Mayr.**B. heeri depilis** Emery.

The smallest of the New England ants. It nests under stones in shady woods and has habits similar to those of *Lasius*. It attends root Coccidæ. The males and winged females make their appearance about the middle of August.

Colebrook (W. M. W.).

Prenolepis Mayr.**P. imparis** Say.

I have not found this ant in the Litchfield Hills. It makes small crater nests in shady oak woods in soil usually containing more or less clay. The workers visit trees for the purpose of attending aphids, obtaining the secretion of extrafloral nectaries, etc. After imbibing these liquids, the gaster often becomes so distended that it is four or five times its normal size and the insects walk with difficulty. In this replete condition *imparis* workers may be said to represent a temporary stage of the more extraordinary enlargement of the gaster seen in the honey ants (*Myrmecocystus*) of the Southwestern States and Mexico. The males and females of *imparis* often pass the winter in the parental nest and celebrate their nuptial flight early in the spring.

New Haven, Yalesville (H. L. V.); Branford (H. W. W.); New Haven (W. E. B.).

P. imparis var. **minuta** Emery.

Differs from the preceding merely in the smaller size of the worker. It is probably not a true variety but merely a nest variation (incipient colony form).

New Haven, Yalesville (H. L. V.).

Subgenus **Nylanderia**.°**P. (N.) parvula** Mayr.

Undoubtedly occurs in southern Connecticut. I have taken it as far east as Mamaroneck and Cold Spring Harbor, N. Y., but have never been able to find it in the Litchfield Hills.

Lasius Fabricius.*Key to Species.*

- | | |
|--|---|
| 1. Maxillary palpi 6-jointed (<i>Lasius s. str.</i>) | 2 |
| Maxillary palpi 3-jointed (subgenus <i>Acanthomyops</i>)..... | 7 |

sively subterranean in its habits. It may often be seen visiting the foliage of trees and bushes in search of aphids and small insects. Professor S. A. Forbes has shown that it is of considerable economic importance on account of its noxious habit of cultivating the root aphids of maize, or Indian corn (*Aphis maidiradicis*). The males and winged females appear in August.

New Haven, West Haven, Branford (H. L. V.); New Haven (W. E. B., B. H. W.); Colebrook, Winsted, Norfolk (W. M. W.).

L. niger Linnæus, var. **neoniger** Emery.

Differs from the preceding variety in having erect hairs on the legs and antennal scapes in the workers and females.

New Hartford, Stafford (W. E. B.); Colebrook, Winsted, Norfolk (W. M. W.).

L. flavus nearcticus Wheeler.

The American representative of the European *flavus*, under which name it is sometimes recorded in the literature. The bodies of the workers have a milky white appearance. The colonies, which are rather small, nest under stones or leaf-mold in damp, shady woods. The males and winged females appear during the first week of August.

Connecticut (Mayr); Colebrook (W. M. W.).

L. brevicornis Emery.

The worker of this species differs from that of the preceding in having the antennal scapes not reaching beyond the posterior corners of the head. The colonies nest under stones on hill slopes and in pastures where the soil is rather dry and sandy. The males and winged females appear about the middle of August.

Branford (H. W. W.); Colebrook (W. M. W.).

L. umbratus mixtus Nylander, var. **aphidicola** Walsh.

Nesting under stones or in old logs and stumps in damp, shady woods. The colonies, which are rather populous, cultivate snow-white root aphids and coccids in great numbers, especially during the winter and early spring. The males and females appear during August and early September.

Westport (W. E. B.); Colebrook, (W. M. W.).

L. umbratus mixtus var. **speculiventris** Emery.

This form, originally described as a distinct species, is scarcely

more than a variety. Its habits, according to my observations, are very similar to those of *aphidicola*.

Colebrook (W. M. W.).

Subgenus *Acanthomyops* Mayr.

L. (A.) interjectus Mayr.

The yellow *Lasii* of the subgenus *Acanthomyops*, besides having only 3- instead of 6-jointed maxillary palpi in the worker and female phases, have a peculiar and rather agreeable odor like lemon verbena, and quite unlike the odor of the typical *Lasii*. They all form large colonies and lead a subterranean aphidicolous existence. *L. interjectus* is the largest species of the genus. It is found nesting in old logs and stumps in open woods and occasionally makes rough mounds or merely excavates its galleries under large stones.

Connecticut (Mayr); Colebrook (W. M. W.).

L. (A.) claviger Roger.

The commonest of our species of *Acanthomyops*, nesting under stones along the edges of woods where there is plenty of warmth and moisture. The males and winged females may be found in the nests from the middle of August till the latter part of September.

Connecticut (Mayr); Colebrook (W. M. W.).

L. (A.) latipes Walsh.

Rather common in grassy fields under stones. It has been shown by Mr. J. F. McClendon and myself that some colonies of this ant have dimorphic females. One of these females (the α -female) is very hairy, and has much flattened femora and tibiae. The other female (the β -female) is intermediate in structure between the β -female and the female of *claviger*. The males and winged females are found in the nest during the latter part of August.

Colebrook (W. M. W., J. F. McClendon).

Formica Latreille.

Key to Species.

- | | |
|--|---|
| 1. Clypeus with a notch in middle of anterior border (<i>F. sanguinea</i>) | 2 |
| Clypeus without a notch in its anterior border | 4 |

females establish their colonies in depauperate colonies of *fusca* var. *subsericea*.

Connecticut (Mayr); Branford, North Haven, New Haven (H. L. V.); New Hartford, Stafford (W. E. B.); Cromwell, Hartford (Forel); Colebrook (W. M. W.).

F. truncicola obscuriventris Mayr.

A single colony, found near the summit of one of the Litchfield Hills (about 1,400 feet).

Connecticut (Mayr); Colebrook (W. M. W.); Brookfield (E. L. Dickerson).

F. truncicola integra Nylander.

Our largest and most conspicuous form of *truncicola* nesting in great colonies which often comprise several nests. These are in piles of large stones or in old logs and stumps. The ants stuff all the crannies of their abodes with bits of dead grass, leaves, etc. Like most other species of *Formica*, *integra* is much given to attending aphids. It is most abundant in hilly regions, where it prefers sunny glades or clearings in the forests. The males and winged females appear in July.

Connecticut (Mayr); Colebrook (W. M. W.).

F. difficilis Emery, var. **consocians** Wheeler.

In this interesting species, as I have shown, the females, which are yellow and hardly larger than the largest workers, are temporary parasites in the nests of *schaufussi* var. *incerta*. Soon after fertilization the queen seeks adoption in some depauperate and probably queenless colony of *incerta* and there permits her hosts to bring up her young. Later the *incerta* workers die off, leaving the *consocians* as a pure and independent colony, which grows rapidly in size and shows no evidence of its parasitic origin. The nesting habits of *difficilis* resemble those of *integra* on a small scale.

Colebrook (W. M. W.).

F. nepticula Wheeler.

Like the preceding, this species has very small females, which, in all probability, are social parasites in the colonies of some other *Formica*, probably *neogagates* Emery. The males and winged females make their appearance during July.

Colebrook (W. M. W.).

F. pallide-fulva schaufussi Mayr.

This is one of the commonest species of *Formica*. It nests in rather small colonies under stones or in small, obscure mound nests in sunny and grassy fields. It is timid and runs rapidly. Its food seems to consist very largely of the excrement of aphids and the carcasses of insects.

Connecticut (Mayr and Emery); New Haven (W. E. B.); Winsted, Norfolk, Colebrook (W. M. W.).

F. pallide-fulva schaufussi var. *incerta* Emery.

Common in the same localities as the typical *schaufussi*, from which it differs merely in somewhat darker coloration and in having fewer hairs on the chin and petiolar border. It is the host of *difficilis* var. *consocians*.

Branford (H. W. W.); Rockville (H. L. V.); Winsted, Norfolk, Colebrook (W. M. W.).

F. pallide-fulva nitidiventris Emery.

The workers are smaller than those of the two preceding forms, dark colored, without hairs on the chin and petiolar border, and with more shining and less pubescent gaster. The habits are similar to those of other forms of the species.

New Haven (P. L. B.); Salisbury, New Haven, Orange (W. E. B.); Colebrook (W. M. W.).

°**F. pallide-fulva nitidiventris** var. *fuscata* Emery.

This variety, which is characterized by its dark color and somewhat opaque gaster, can hardly be absent from Connecticut, as it occurs in the adjacent states.

F. fusca Linnæus, var. *subsericea* Say. Silky Ant.

Next to *Lasius niger* var. *americanus*, this is the commonest of our ants and hence also of our insects. It prefers sunny, grassy places, and either constructs dome-shaped mounds which are largest and most definite in outline in the Middle States, or excavates its galleries under stones, boards, the bark of stumps, etc. Except when living in large colonies, it is a very cowardly species. Like the other members of the genus *Formica*, it attends aphids, but is equally fond of feeding on the dead bodies of insects. The males and winged females make their appearance during July and August.

Suffield (Dimmock); Branford, Cheshire, Mt. Carmel, New Haven (H. L. V.); New Haven, Salisbury (W. E. B.); Cromwell, Hartford (Forel); Winsted, Norfolk, Colebrook (W. M. W.).

F. fusca var. *subaenescens* Emery.

A rare species, apparently, in New York and New England, but common in the Northern Middle States (Illinois, Wisconsin, Michigan). It differs from the preceding variety in having a more metallic and less pubescent surface. It prefers to nest under logs and stones in rather shady woods.

Connecticut (Emery); Colebrook (W. M. W.).

°**F. subpolita** Mayr.

I have not seen specimens of the typical form of this species from the State. It is possible that Mayr's specimens may have belonged to the following species.

Connecticut (Mayr).

F. neogagates Emery.

Nesting in rather small colonies under stones only on the hills at an altitude of about 1,000 feet or more, according to my observations. The males and winged females appear during late July and early August.

Kent, Salisbury (W. E. B.); Norfolk, Colebrook (W. M. W.).

Polyergus Latreille.

P. lucidus Mayr.

This rare and beautiful species, the "shining slave-maker" of McCook, or "shining amazon," as it may be called, uses the workers of *Formica schaufussi* as slaves, or auxiliaries. These are bred from pupæ kidnapped from their maternal nests by the warlike *lucidus* workers. The latter are quite unable to feed themselves, excavate their nests, or care for their own brood, but have to depend for these important activities on the *schaufussi* workers. Hence the ants of this species are quite unable to live an independent life and may be regarded as permanently parasitic on fragments of *schaufussi* colonies which they bring together with great skill. The sexual forms make their appearance during August.

Connecticut (Mayr).

Camponotus Mayr.

Key to Species.

1. Clypeus with a distinct notch or impression in the middle of its anterior border **fallax** and its varieties
Clypeus without such a notch or impression..... 2
2. Head of worker major smooth and shining behind; color, at least in part, light red or yellow (**castaneus**) 3
Head of worker major opaque or feebly shining behind; color black, or black and dark red (**herculeanus**) 4
3. Yellow or light red; gaster slightly darker (typical) **castaneus**
Head black or dark brown **castaneus** subspecies **americanus**
4. Gaster opaque, with long, appressed pubescence 5
Gaster shining, with short, sparse pubescence; thorax deep red ... **herculeanus** subspecies **ligniperda** var. **noveboracensis**
5. Deep black throughout ... **herculeanus** subspecies **pennsylvanicus**
Legs, posterior portion of thorax, petiole, and base of gaster brownish red **herculeanus** subspecies **pennsylvanicus** var. **ferrugineus**

C. fallax Nylander, var. **nearcticus** Emery.

Till recently this species has been cited in the literature as *C. marginatus* Latreille. Our American subspecies and varieties nest in the hollow twigs of trees and bushes and attend aphids.

Connecticut (Mayr); Colebrook (W. M. W.).

C. castaneus Latreille.

The typical form of this species is probably confined to the lower, warmer, and southernmost portions of the State, as I have seen no trace of it in the Litchfield Hills. It nests under stones and logs in rather small colonies.

Connecticut (Mayr, Coe); Westville (W. E. B.).

C. castaneus americanus Mayr.

Brookfield (E. L. Dickerson).

C. herculeanus pennsylvanicus Degeer. Carpenter Ant.

The common "carpenter ant," entirely black in color. It nests usually in shady woods in old logs and stumps, whence it may migrate into old farm-houses and suburban residences, and become a pest, both by riddling the wood-work with its large anastomosing galleries and by visiting the pantries and kitchens for sweets.

Connecticut (Mayr); Woodmont (P. L. B.); New Haven, Branford (H. L. V.); Colebrook (W. M. W.).

C. herculeanus pennsylvanicus var. **ferrugineus** Fabricius.

A beautiful color-variety of *pennsylvanicus*, with the legs, inferior and posterior portions of the thorax, petiole, and base of gaster rust-red in the female and worker phases. Its habits are very similar to those of the typical form, but it seems to be less abundant and more local in its distribution. I have been unable to find it in the Litchfield Hills.

New Haven (E. J. S. M., H. L. V.); Orange, New Canaan (W. E. B.).

C. herculeanus ligniperda var. **noveboracensis** Fitch.

Nesting in old stumps and logs like the preceding, from which it differs in having a smoother surface and an entirely red thorax in the worker phases.

New Hartford, Orange (W. E. B.); Colebrook (W. M. W.).